

2009

65. Bielak, L. F., Horenstein, R. B., Ryan, K. A., et al. (2009). Circulating CD34+ cell count is associated with extent of subclinical atherosclerosis in asymptomatic Amish men, independent of 10-year Framingham risk. *Clin Med Cardiol* 3: 53-60.
66. Cheng, Y. C., Kao, W. H., Mitchell, B. D., et al. (2009). Genome-wide association scan identifies variants near matrix metalloproteinase (MMP) genes on chromosome 11q21-22 strongly associated with serum MMP-1 levels³. *Circ Cardiovasc Genet* 2(4): 329-337.
67. Daley, E., Streeten, E. A., Sorkin, J. D., et al. (2009). Variable bone fragility associated with an Amish COL1A2 variant and a knock-in mouse model. *J Bone Miner Res* 25(2): 247-261.
68. Heard-Costa, N. L., Zillikens, M. C., Monda, K. L., et al. (2009). NRXN3 is a novel locus for waist circumference: a genome-wide association study from the CHARGE Consortium. *PLoS Genet* 5(6): e1000539.
69. Michos, E. D., Streeten, E. A., Ryan, K. A., et al. (2009). Serum 25-hydroxyvitamin D levels are not associated with subclinical vascular disease or C-reactive protein in the Old Order Amish. *Calcif Tissue Int* 84(3): 195-202.
70. Naj, A. C., Kao, W. H., O'Connell, J. R., Mitchell, B. D. and Silver, K. D. (2009). Sequence variation in IGF1R is associated with differences in insulin levels in nondiabetic Old Order Amish. *Diabetes Metab Res Rev* 25(8): 773-779.
71. Perry, J. R. B., Stolk, L., Franceschini, N., et al. (2009). Meta-analysis of genome-wide association data identifies two loci influencing age at menarche. *Nature Genetics* 41(6): 648-650.
72. Prokopenko, I., Zeggini, E., Hanson, R. L., et al. (2009). Linkage disequilibrium mapping of the replicated type 2 diabetes linkage signal on chromosome 1q. *Diabetes* 58(7): 1704-1709.
73. Shen, H., Herzog, W., Drolet, M., et al. (2009). Aspirin resistance in healthy drug-naive men versus women (from the Heredity and Phenotype Intervention Heart Study). *Am J Cardiol* 104(4): 606-612.
74. Shen, H., Pollin, T. I., Damcott, C. M., et al. (2009). Glucokinase regulatory protein (GCKR) gene polymorphism affects postprandial lipemic response in a dietary intervention study. *Hum Genet* 126: 567-574.
75. Shuldiner, A. R., O'Connell, J. R., Bliden, K. P., et al. (2009). Association of cytochrome P450 2C19 genotype with the antiplatelet effect and clinical efficacy of clopidogrel therapy. *JAMA* 302(8): 849-857.
76. Tarasov, K. V., Sanna, S., Scuteri, A., et al. (2009). COL4A1 is associated with arterial stiffness by genome-wide association scan. *Circ Cardiovasc Genet* 2(2): 151-158.
77. Wang, Y., O'Connell, J. R., McArdle, P. F., et al. (2009). Whole-genome association study identifies STK39 as a hypertension susceptibility gene. *Proc Natl Acad Sci USA* 106(1): 226-231.