

ORIGINAL ARTICLE



Efficacy and Safety of an mRNA Seasonal Influenza Vaccine in Adults

Authors: Isabel Leroux-Roels, M.D., Grace Huang, M.D., Murdo Ferguson, M.D., Anita Kohli, M.D., Rebecca Clark, M.D., Markus Bickel, M.D., Mieke Soens, M.D., ⁺⁶, for the Fluent Trial Investigators* [Author Info & Affiliations](#)

Published May 6, 2026 N Engl J Med 2026;394:1803-1813 DOI: 10.1056/NEJMoa2516491 VOL. 394 NO. 18

Copyright © 2026



Abstract

BACKGROUND

Seasonal influenza causes substantial illness and death in adults 50 years of age or older, even with current vaccines. An investigational messenger RNA (mRNA)–based vaccine called mRNA-1010 encodes hemagglutinin glycoproteins from World Health Organization–recommended influenza strains.

METHODS

In this phase 3, double-blind, active-controlled trial, we randomly assigned adults 50 years of age or older to receive trivalent mRNA-1010 (37.5 µg, which includes 12.5 µg of each strain) or a licensed standard-dose comparator. The primary efficacy end point was relative vaccine efficacy against reverse-transcriptase–polymerase-chain-reaction (RT-PCR)–confirmed, protocol-defined influenza-like illness caused by influenza A or B, from at least 14 days after vaccination through the end of the influenza season. Hypothesis testing was conducted hierarchically to assess noninferiority (lower boundary of the 95% confidence interval [CI], >–10%), superiority (lower boundary of the 95% CI, >0%), and a higher level of superiority (lower boundary of the 95% CI, >9.1%).

RESEARCH SUMMARY




Efficacy and Safety of an mRNA Seasonal Influenza Vaccine in Adults

RESULTS

A total of 40,703 participants received mRNA-1010 (20,350 participants) or the standard-dose comparator (20,353 participants); the median follow-up was 181 days (range, 1 to 227). RT-PCR–confirmed, protocol-defined influenza-like illness was observed in 411 of 20,170 recipients of mRNA-1010 (2.0%) and 1,557 of 20,124 recipients of the standard-dose comparator (2.8%), which corresponded to a relative vaccine efficacy of 29.8% (95% CI, 15.7 to 35.4), thereby meeting the criteria for noninferiority, superiority, and a higher level of superiority. Adverse reactions were more frequent with mRNA-1010 than with the standard-dose comparator (injection-site pain in 65.8% vs. 29.8%, fatigue in 45.1% vs. 20.3%, headache in 37.8% vs. 18.0%, and myalgia in 35.4% vs. 11.6%); most reactions were mild to moderate and

In this trial, mRNA-1010 was superior to standard-dose licensed vaccines for prevention of RT-PCR–confirmed, protocol-defined influenza-like illness in adults 50 years of age or older. Solicited adverse reactions were more frequent with mRNA-1010. (Funded by Blackstone Life Sciences and Moderna; Fluent ClinicalTrials.gov number, [NCT06602024](#).)

QUICK TAKE**An mRNA Seasonal Influenza Vaccine for Adults**

2m 34s

Seasonal influenza viruses are estimated by the World Health Organization (WHO) to cause 3 to 5 million cases of severe illness and up to 650,000 deaths annually, posing a persistent global public health challenge.¹ Improved vaccine efficacy in adults 50 years of age or older is needed, given the increased morbidity and mortality in this population.^{2,3}

Enhanced (high-dose, adjuvanted, or recombinant-based) influenza vaccines provide greater protection in older adults than standard-dose formulations, although effectiveness varies according to season and match to circulating strains.^{4,5} Most licensed vaccines rely on egg-based platforms.⁶ Egg-based production is time-consuming and can introduce egg-adaptive mutations, which reduce antigenic match with circulating strains.⁶ A need remains for influenza vaccines that can be rapidly updated to evolving strains and elicit robust protection.

The investigational seasonal influenza vaccine mRNA-1010 encodes surface hemagglutinin (HA) antigens from the three WHO-recommended 2024–2025 strains for cell- or recombinant-based vaccines: A/H1N1, A/H3N2, and B/Victoria.⁷ By avoiding egg-based production and enabling rapid strain updates, the messenger RNA (mRNA) platform addresses key limitations of current influenza vaccines and may improve protection in older adults.⁸

The Fluent trial was conducted during the 2024–2025 Northern Hemisphere influenza season to evaluate the relative vaccine efficacy of mRNA-1010 as compared with licensed standard-dose influenza vaccines in preventing laboratory-confirmed influenza-like illness in adults 50 years of age or older. Here, we report the findings from the primary analyses of this pivotal phase 3 trial.

Methods

TRIAL DESIGN AND OVERSIGHT

This phase 3, double-blind, randomized, active-controlled trial enrolled participants at 301 sites in 11 countries in the Northern Hemisphere. The trial was approved by appropriate institutional review boards and was conducted according to the applicable principles of the International Council for Harmonisation Good Clinical Practice guidelines, the E6(R2) Good Clinical Practice guidelines, the principles of the Declaration of Helsinki, and all local laws or regulations. All the participants provided written informed consent before enrollment. Oversight was provided by an independent data and safety monitoring board.

A sponsor, Moderna, was responsible for overall trial design, site selection, monitoring, and data analysis, which were facilitated by Parexel International. The authors vouch for the accuracy and completeness of the data and for the fidelity of the trial to the protocol and statistical analysis plan. The manuscript was prepared by employees of Moderna with assistance from medical writers funded by Moderna. Additional information is provided in the [Supplementary Appendix, protocol](#), and statistical analysis plan, all available with the full text of this article at NEJM.org.

TRIAL PARTICIPANTS

Eligible participants were adult [Have questions about this content? Try AI Companion.](#) [NEW](#) [dition](#). A complete list of eligibility criteria is provided in the [Supplementary Appendix, protocol](#).

Tetra, Influsplit Tetra, or Alpharix Tetra) with the use of a centralized interactive response technology system that implemented the prespecified randomization schedule and concealed treatment allocation until assignment. A licensed trivalent standard-dose comparator was preferred; a licensed quadrivalent comparator was used where a trivalent vaccine was not available. In countries recommending higher-dose or adjuvanted influenza vaccines in older adults, potential participants were informed of this recommendation. Vaccines were shipped frozen and stored at clinical sites before preparation and injection (see the [Supplementary Appendix](#)).

EFFICACY ASSESSMENTS

The primary efficacy objective was to evaluate the relative vaccine efficacy of mRNA-1010 as compared with a standard-dose comparator against the first episode of protocol-defined influenza-like illness caused by any influenza A or B strain confirmed by a reverse-transcriptase–polymerase-chain-reaction (RT-PCR) assay, beginning at least 14 days after injection through the end of the influenza season. Efficacy end-point events were identified through active surveillance, including twice-weekly e-diary prompts to report respiratory symptoms from day 1 through the end of the influenza season.

If a participant met criteria for protocol-defined respiratory illness, a nasopharyngeal swab was collected within 72 hours after symptom onset and tested for influenza with an RT-PCR assay (see the [Supplementary Appendix](#)). Protocol-defined influenza-like illness was an influenza infection confirmed by RT-PCR assay and the presence of at least one systemic symptom (temperature $>37.2^{\circ}\text{C}$, chills, feverish, tiredness, headaches, or myalgia) and at least one respiratory symptom (sore throat, cough, sputum production, wheezing, or difficulty breathing) within 7 days before or after the positive test. Key secondary objectives were to evaluate the relative vaccine efficacy of mRNA-1010 as compared with the standard-dose comparator against RT-PCR–confirmed, modified Centers for Disease Control and Prevention (CDC)–defined influenza-like illness (temperature $>37.2^{\circ}\text{C}$ and cough or sore throat⁹) and against protocol-defined influenza-like illness with strains with antigenic match to the vaccine strains.

SAFETY ASSESSMENTS

The primary safety objective was to evaluate the safety and reactogenicity of mRNA-1010. Unsolicited adverse events were assessed through 28 days after injection with the use of active and passive surveillance. Data on adverse events leading to discontinuation of trial participation, medically attended adverse events, serious adverse events, and adverse events of special interest were collected through day 181 or the data cut-off date (April 30, 2025). Unsolicited adverse events were graded as mild, moderate, or severe. Data on solicited local and systemic adverse reactions were collected from approximately 6000 participants who used an e-diary for the first 7 days after injection (see the [Supplementary Appendix](#)).

STATISTICAL ANALYSIS

The planned sample size was approximately 56,000 participants undergoing randomization over two influenza seasons, with approximately 70% enrolled during the first year. Accrual of 836 cases in the per-protocol efficacy analysis population was estimated to provide approximately 80% power to show noninferiority with a noninferiority margin of 10%, under the assumption of a true relative vaccine efficacy of 10% at an overall one-sided type I error rate of 2.5% (additional details are provided in the [Supplementary Appendix](#)). Efficacy analyses were prespecified at the end of both seasons, with the Pocock spending function used to control the overall type I error rate. A stratified Cox proportional-hazards model was used to estimate the hazard ratio. Relative vaccine efficacy was calculated as $(1 - \text{hazard ratio [mRNA-1010 vs. standard-dose comparator]}) \times 100\%$, with stratification according to age (50 to 64 or ≥ 65 years) and status with respect to influenza vaccination during the previous influenza season. Missing efficacy data were not imputed.

The primary objective was assessed sequentially for noninferiority (lower boundary of the 95% confidence interval [CI] of the relative vaccine efficacy, $>-10\%$), superiority (lower boundary of the 95% CI, $>0\%$), and higher-level superiority (lower boundary of the 95% CI, $>9.1\%$), on the basis of correspondence with the Food and Drug Administration; these values corresponded to hazard ratios of 1.10, 1.05, and 1.01, respectively. Noninferiority was concluded if the one-sided P value was <0.025 . Have questions about this content? Try AI Companion. Superiority and higher-level superiority were tested sequentially with the use of the same alpha-spending framework (null hypothesis: hazard ratio, ≥ 1.00 and ≥ 0.909 , respectively). NEW

A/H3N2, and B/Victoria was 29.6% (95% CI, 16.4 to 40.7), 22.2% (95% CI, 4.3 to 36.9), and 29.1% (95% CI, -18.5 to 57.5), respectively (Table 2).

The favorable relative vaccine efficacy of mRNA-1010 over the standard-dose comparator also appeared evident across subgroups defined according to age, sex, race, BMI, comparator vaccine (trivalent or quadrivalent), geographic region, baseline risk status, frailty, and status with respect to influenza vaccination during the previous influenza seasons (Table S3). The estimates of relative vaccine efficacy were generally consistent for trivalent and quadrivalent comparator vaccines, with overlapping confidence intervals. Among the 19,260 participants 65 years of age or older, the relative vaccine efficacy was 27.4% (95% CI, 12.1 to 40.0). The estimates of relative vaccine efficacy were also consistent in high-risk persons (Figure 1). An analysis of the cumulative number of cases over time suggested that the efficacy advantage for mRNA-1010 was observed early and maintained throughout the full influenza season (Figure 2).

In exploratory analyses, medically attended influenza-associated outcomes across all health care settings occurred in 80 recipients of mRNA-1010 and 120 recipients of the standard-dose comparator, which corresponded to a relative vaccine efficacy of 33.7% (95% CI, 12.0 to 50.0) (Fig. S2). Among 64 participants who sought higher-level care (hospitalization, emergency department, or urgent care), health care encounters occurred in 22 recipients of mRNA-1010 and 42 recipients of the standard-dose comparator, which corresponded to a relative vaccine efficacy of 47.9% (95% CI, 12.8 to 68.9). The number of participants who were hospitalized was lower with mRNA-1010 than with the standard-dose comparator (4 vs. 8), as was the number of participants who visited the emergency department (6 vs. 12), findings consistent with the overall relative vaccine efficacy for higher-level care.

SAFETY

Solicited local and systemic adverse reactions were reported more frequently with mRNA-1010 than with the standard-dose comparator; most reactions were mild to moderate and transient (median duration, 1 to 2 days) (Figure 3). Injection-site pain was the most common solicited local adverse reaction both with mRNA-1010 and with the standard-dose comparator (65.8% vs. 29.8%); the most frequent solicited systemic adverse reactions were fatigue (45.1% vs. 20.3%), headache (37.8% vs. 18.0%), and myalgia (35.4% vs. 11.6%). Grade 3 solicited adverse reactions were reported by 6.4% of recipients of mRNA-1010 and 1.0% of recipients of the standard-dose comparator (Table S4). No grade 4 solicited adverse reactions were reported. The frequency of solicited adverse reactions was lower among older participants than among younger participants (Table S5).

The frequency of unsolicited adverse events, including serious adverse events, adverse events leading to discontinuation of trial participation, adverse events of special interest, was similar in the two groups (Tables S6 and S7, respectively). Have questions about this content? Try AI Companion. vaccine-related unsolicited adverse events through the data-cutoff date occurred more frequently with mRNA-1010 than with the standard-dose comparator (0.5% [104 of 20,350

Confirmed, protocol-defined influenza-like illness caused by any influenza A or B strain			
Caused by any influenza A	386 (1.9)	522 (2.6)	26.5 (16.1 to 35.5)§
Caused by influenza A/H1N1	223 (1.1)	315 (1.6)	29.6 (16.4 to 40.7)§
Caused by influenza A/H3N2	158 (0.8)	202 (1.0)	22.2 (4.3 to 36.9)§
Caused by influenza B	25 (0.1)	35 (0.2)	29.1 (-18.5 to 57.5)§
First occurrence of RT-PCR-confirmed, modified CDC-defined influenza-like illness caused by any influenza A or B strain	223 (1.1)	290 (1.4)	23.5 (9.0 to 35.8)†‡
Caused by any influenza A	211 (1.0)	276 (1.4)	24.0 (9.1 to 36.5)§
Caused by influenza A/H1N1	125 (0.6)	172 (0.9)	27.7 (9.0 to 42.6)§
Caused by influenza A/H3N2	83 (0.4)	102 (0.5)	19.1 (-8.1 to 39.5)§
Caused by influenza B	12 (<0.1)	14 (<0.1)	14.8 (-84.3 to 60.6)§

* Data are from the per-protocol efficacy analysis population, which included participants who had undergone randomization, received a trial vaccine, and had no major protocol deviations affecting the efficacy outcomes as determined before database lock and unblinding. The event is the first reverse-transcriptase-polymerase-chain-reaction (RT-PCR)-confirmed, protocol-defined influenza-like illness or first RT-PCR-confirmed, modified Centers for Disease Control and Prevention (CDC)-defined influenza-like illness that begins at least 14 days after trial vaccination through the end of influenza season caused by any influenza A or B strain, regardless of vaccine match. Data for participants without the event are censored at early discontinuation of trial participation or death unrelated to influenza; the date of early RT-PCR-confirmed, protocol-defined influenza-like illness starting fewer than 14 days after trial vaccination; the end of the trial; the end of influenza season; or the data-cutoff date, whichever occurs the earliest. Relative vaccine efficacy is defined as (1-hazard ratio [mRNA-1010 vs. standard-dose comparator]) × 100. The relative vaccine efficacy and the confidence interval are based on a stratified Cox proportional-hazards model with vaccine group as a fixed effect, with adjustment for the randomization stratification factors: age group (50 to <65 years or ≥65 years) and the status with respect to influenza vaccination during the previous influenza season (received or not received). Efron's method was used to handle ties.

† Hypothesis testing was conducted for noninferiority, superiority, and higher-level superiority in a hierarchical fashion at a one-sided significance level of 2.5%. The success criteria were equivalent to a lower boundary of the 95% confidence interval of the relative vaccine efficacy of more than -10% for noninferiority, more than 0% for superiority, and more than 9.1% for higher-level superiority.


‡ One-sided P<0.001 for noninferiority, superiority, and higher-level superiority.

§ No hypothesis testing was conducted; the 95% confidence interval was provided as a descriptive measure. The width of the confidence interval was not adjusted for multiplicity and should not be used for hypothesis testing.

¶ One-sided P<0.001 for noninferiority; one-sided P=0.001 for superiority; and one-sided P=0.03 for higher-level superiority.


Relative Vaccine Efficacy (Per-Protocol Efficacy Population).

FIGURE 1



Relative Vaccine Efficacy of mRNA-1010 as Compared with Licensed Influenza Vaccines According to High-Risk Subgroup (Per-Protocol Efficacy Population).

FIGURE 2




First Occurrence of Protocol-Defined Influenza-like Illness Caused by Any Influenza A or B Strain.

FIGURE 3



and 1.5% of recipients of the standard-dose comparator through the data-cutoff date (Table S7) and were most frequently classified under the Medical Dictionary for Regulatory Activities system organ class of infections and infestations (0.4% in each group) (Table S8). Three recipients of mRNA-1010 (<0.1%) and two recipients of the standard-dose comparator (<0.1%) had serious adverse events considered by the investigator to be vaccine-related (Tables S7 and S9). Fatal events occurred in 0.2% of the participants in each group; none were considered by the investigator to be vaccine-related.

Protocol-defined adverse events of special interest were reported in 15 participants (<0.1%) in each group through the data-cutoff date. A total of 4 participants (2 per group) had adverse events of special interest that were considered by the investigator to be vaccine-related, one of which occurred within 28 days after vaccination (Bell's palsy in a recipient of the standard-dose comparator). Adverse events of special interest in recipients of mRNA-1010 included thrombocytopenia 84 days after vaccination and cardiomyopathy 95 days after vaccination, which was not considered by the independent Cardiac Event Adjudication Committee (CEAC) to be an event of myopericarditis. One recipient of the standard-dose comparator had pericarditis 60 days after vaccination. As assessed by the CEAC, there were no cases of acute myocarditis, myopericarditis, or pericarditis among recipients of mRNA-1010 within the conservative risk window of 42 days after vaccination.



Solicited Local and Systemic Reactions.

Discussion

The investigational vaccine mRNA-1010 is based on a proprietary mRNA platform designed to prevent influenza disease caused by influenza A or B strains. Other mRNA vaccines on this platform have shown clinical success against respiratory tract infections.^{10,11} The effectiveness of influenza vaccines varies according to season and circulating strains. For the 2024–2025 season, efficacy estimates of licensed, predominately egg-based influenza vaccines ranged from 36 to 54% in the United States and Europe.^{12,13} The mRNA platform enables high-fidelity antigen encoding, avoids egg-based production, and can be rapidly updated to minimize antigen mismatch and manufacturing time,^{14,15} which may contribute to improved vaccine efficacy.

In this international, phase 3 trial involving adults 50 years of age or older, a single dose of mRNA-1010 met the prespecified criterion of higher-level superiority relative to a licensed standard-dose comparator, with an estimated relative vaccine efficacy of 26.6% (95% CI, 16.7 to 35.4). The superior relative vaccine efficacy of mRNA-1010 was observed early and maintained throughout the full influenza season. In the high-transmission 2024–2025 Northern Hemisphere influenza season, the efficacy advantage for mRNA-1010 was observed across influenza strains. Similar estimates of relative vaccine efficacy were observed across all vaccine-included strains: A/H1N1 (29.6%), A/H3N2 (22.2%), and B/Victoria (29.1%). The mRNA platform enables precise HA sequence matching to circulating strains, reflected in consistent efficacy across subtypes.

Among more than 19,000 adults 65 years of age or older, a population particularly susceptible to severe influenza, the relative vaccine efficacy of mRNA-1010 was 27.4% (95% CI, 12.1 to 40.0), which suggests benefits similar to those of enhanced (high-dose, adjuvanted, or recombinant-based) influenza vaccines in older adults.¹⁶ Similar efficacy was observed in other prespecified high-risk subgroups, including participants with obesity or underlying high-risk medical conditions.

In exploratory analyses, recipients of mRNA-1010 had fewer cases of medically attended influenza illness across all levels of health care use than recipients of the standard-dose comparator. These results align with those of previous studies showing reduced influenza-like illness–associated hospitalizations with high-dose influenza vaccines as compared with standard-dose formulations.^{17,22} Although the trial was not powered for these outcomes, the consistency of relative vaccine efficacy across these many outcomes suggests that mRNA-1010 efficacy is broadly applicable across diverse populations and the spectrum of influenza disease.

This trial had several strengths and limitations. Its global design and inclusion of demographically and clinically diverse participants support the generalizability of the findings. Although case numbers were limited in some subgroups (including participants ≥ 75 years of age and vulnerable or frail adults), which resulted in wide confidence intervals, the magnitude of the point estimate of relative vaccine efficacy was similar to that for the primary end point. Although the analysis covered only one influenza season, it included 968 confirmed cases across all three circulating strains. The number of influenza B cases was small (60 cases); however, the point estimate of relative vaccine efficacy was consistent with overall relative vaccine efficacy and aligned with previous data showing superior immune responses against B lineages with mRNA-1010.²³ Because persons 65 years of age or older are at increased risk for severe influenza, some countries recommend enhanced influenza vaccines. However, this multinational Northern Hemisphere trial was conducted across regions with differing standards of care; therefore, a standard-dose comparator was selected, and this information was included in the informed consent. The magnitude of the relative vaccine efficacy observed with mRNA-1010 relative to the standard-dose comparator appears to be consistent with that of other enhanced vaccines¹⁶ and consistent with previous results in which mRNA-1010 showed superior immunogenicity as compared with a licensed standard-dose and high-dose comparator in older adults.²³

In this trial, vaccination with mRNA-1010 was superior to a licensed standard-dose comparator in preventing influenza. Solicited adverse reactions were more common in the mRNA-1010 group, which suggests a need for balancing assessment of temporary vaccine-induced reactogenicity events with the magnitude of protection against influenza. Serious adverse events were infrequent and similar in the two groups. These findings support the role of mRNA-1010 in improving influenza prevention.

NOTES

A [data sharing statement](#) provided by the authors is available with the full text of this article at NEJM.org.

Supported by Blackstone Life Sciences and Moderna.

[Disclosure forms](#) provided by the authors are available with the full text of this article at NEJM.org.

We thank the participants in the trial and their families; the members of the data and safety monitoring board for their hard work, support, and guidance of the trial; Agi Buchanan, M.D., Ph.D., Andrei Avanesov, Ph.D., Ren Chen, Ph.D., Haritha Singireddy, M.S., Brianna Fidler, M.A., Elissa Malkin, D.O., M.P.H., Chelsea Canan, Ph.D., M.P.H., and Peg Mutty, B.S., of Moderna, for helpful discussions; Alana Simorellis, Ph.D., of Moderna, for writing and editorial support with an earlier version of the manuscript; and Aliscia Daniels, Ph.D., of MEDiSTRAVA, for medical writing and editorial assistance with an earlier version of the manuscript, in accordance with Good Publication Practice (GPP3) guidelines, funded by Moderna, and under the direction of the authors.

SUPPLEMENTARY MATERIAL

Protocol (nejmoa2516491_protocol.pdf)

DOWNLOAD 4.71 MB

Supplementary Appendix (nejmoa2516491_appendix.pdf)

DOWNLOAD 931.02 KB

Disclosure Forms (nejmoa2516491_disclosures.pdf)

DOWNLOAD 1.29 MB

Data Sharing Statement (nejmoa2516491_data_sharing_statement.pdf) Have questions about this content? [Try AI Companion.](#)

DOWNLOAD 72.65 KB

NEW

ADVERTISEMENT

[← Go to Citation](#) [Google Scholar](#) [find it](#)

2. Divo MJ, Martinez CH, Mannino DM. Ageing and the epidemiology of multimorbidity. *Eur Respir J* 2014;44:1055-1068.

[← Go to Citation](#) [Crossref](#) [PubMed](#) [Web of Science](#) [Google Scholar](#) [find it](#)

3. Goronzy JJ, Weyand CM. Understanding immunosenescence to improve responses to vaccines. *Nat Immunol* 2013;14:428-436.

[← Go to Citation](#) [Crossref](#) [PubMed](#) [Web of Science](#) [Google Scholar](#) [find it](#)

4. Chaves SS, Naeger S, Lounaci K, et al. High-dose influenza vaccine is associated with reduced mortality among older adults with breakthrough influenza even when there is poor vaccine-strain match. *Clin Infect Dis* 2023;77:1032-1042.

[← Go to Citation](#) [Crossref](#) [PubMed](#) [Web of Science](#) [Google Scholar](#) [find it](#)

5. Youhanna J, Tran V, Hyer R, Domnich A. Immunogenicity of enhanced influenza vaccines against mismatched influenza strains in older adults: a review of randomized controlled trials. *Influenza Other Respir Viruses* 2024;18(4):e13286-e13286.

[← Go to Citation](#) [Crossref](#) [PubMed](#) [Web of Science](#) [Google Scholar](#) [find it](#)

SHOW ALL REFERENCES

ADVERTISEMENT



PHYSICIAN JOBS

MAY 28, 2026

Hematology / Oncology Bay Shore, New York
[Hematologist and Genitourinary Medical Oncologist - Bay Shore, New York](#)

Pulmonary / Critical Care New York
[Pulmonary, Critical Care, and Sleep Opportunities at Northwell Health!](#)

Pediatrics, General New York City, New York
[Southwestern New York State / Adding 2 Pediatricians to Group / Visa Support](#)

Have questions about this content? [Try AI Companion.](#)

NEW

Internal Medicine Texas
[Internal Medicine Physician, \\$280k/yr](#)



MORE LIKE THIS

[ORIGINAL ARTICLE](#) | NEJM | NOV 20, 2025

Efficacy, Immunogenicity, and Safety of Modified mRNA Influenza Vaccine

D. Fitz-Patrick and Others

[ORIGINAL ARTICLE](#) | NEJM | DEC 11, 2025

High-Dose Influenza Vaccine to Reduce Hospitalizations

J. Pardo-Seco and Others

[ORIGINAL ARTICLE](#) | NEJM | DEC 11, 2025

High-Dose Influenza Vaccine Effectiveness against Hospitalization in Older Adults

N.D. Johansen and Others

[ORIGINAL ARTICLE](#) | NEJM | DEC 14, 2023

Recombinant or Standard-Dose Influenza Vaccine in Adults under 65 Years of Age

A. Hsiao and Others

[CLINICAL IMPLICATIONS OF BASIC RESEARCH](#) | NEJM | MAR 23, 2023

An mRNA Influenza Vaccine — Could It Deliver?

K.M. Neuzil

Have questions about this content? [Try AI Companion.](#)

NEW

BASE CATALYSIS OF H2O2 AND H2O2 PROMOTERS FOR SEVERE CORONARY DISEASE

A.O. Gupta and Others

ORIGINAL ARTICLE | MAY 07, 2026 |

Left Ventricular Unloading in High-Risk Percutaneous Coronary Intervention

D. Perera and Others

ORIGINAL ARTICLE | MAY 07, 2026 | CME |


Combined Oral Ivermectin and 5% Permethrin Cream to Treat Severe Scabies

C. Bernigaud and Others

Tap into groundbreaking research and clinically relevant insights

SUBSCRIBE

Already a subscriber? [Activate](#) your online access.



ARTICLE CATEGORIES

- Research
- Reviews
- Clinical Cases
- Perspective
- Commentary
- Other
- Browse all Articles
- Current Issue
- Issue Index

RESOURCES

- Authors & Reviewers
- Submit a Manuscript
- Subscribers
- Institutional Administrators
- Media
- Advertisers
- Agents
- Permissions & Licensing
- Reprints
- NEJM CareerCenter

ABOUT US

- About NEJM
- NEJM Group
- Products & Services
- Editors & Publishers

SUBSCRIPTIONS

- Subscribe
- Private Subscription
- Create Account

Have questions about this content? **Try AI Companion.**


NEW

Help
Site Feedback

STAY CONNECTED

Email Alerts
Create Account
Apps
NEJM CareerCenter
Podcasts
RSS Feed
Remote Access

FOLLOW US

 Facebook
 X
 Instagram
 Youtube
 LinkedIn
 Bluesky
 TikTok
 Threads

PUBLICATIONS

[The New England Journal of Medicine](#)

[NEJM Clinician](#)

[NEJM Evidence](#)

[NEJM AI](#)

[NEJM Catalyst Innovations in Care Delivery](#)

Copyright © 2026 [Massachusetts Medical Society](#). All rights reserved, including those for text and data mining, AI training, and similar technologies. Electronic ISSN 1533-4406. Print ISSN 0028-4793. The content of this site is intended for health care professionals.
[Copyright](#) | [Terms](#) | [Privacy Policy](#) | [Cookie Preferences](#)



Have questions about this content? [Try AI Companion.](#)

NEW